

Sound modes at the BCS-BEC crossover

H. Heiselberg*

Danish Defense Research Establishment, Ryvangsalle' 1, DK-2100 Copenhagen Ø, Denmark

First and second sound speeds are calculated for a uniform superfluid gas of fermi atoms as a function of temperature, density and interaction strength. The second sound speed is of particular interest as it is a clear signal of a superfluid component and it determines the critical temperature. The sound modes and their dependence on density, scattering length and temperature are calculated in the BCS, molecular BEC and unitarity limits and a smooth crossover is extrapolated. It is found that first and second sound undergo avoided crossing on the BEC side due to mixing. Consequently, they are detectable at crossover both as density and thermal waves in traps.

PACS numbers: 03.75.Ss, 67.40.Pm, 05.30.Fk, 05.30.Jp

I. INTRODUCTION

Recent experiments probe systems of strongly interacting Fermi and Bose atoms and molecular bosons at low temperatures. Evidence for a continuous crossover between BCS to a molecular BEC superfluid is found by measurements of expansion [1, 2, 3, 4, 5], collective modes [6, 7], RF spectroscopy [8], thermal energies and entropies [9], and correlations [10]. The next generation of experiments will measure sound velocities in these systems. Second sound has been studied in He^3 , He^4 and BCS superfluids (see, e.g., [11],[12]). Second sound is particularly interesting as it is a clear signal of a superfluid component and determines the critical temperature [13]. The dependence of the sound speeds on temperature, density and interaction strength will constrain the equation of state and models.

Eagles [14] and Leggett [15] described the smooth crossover at zero temperature by a mean field gap equation. Extensions to finite temperature especially around the superfluid critical temperature can be found in [16, 17, 18, 19, 20, 21, 22, 23]. The various crossover models differ in the inclusion of selfenergies, whether one or two channels are employed, etc. Recently, the crossover has been calculated in detail by quantum Monte Carlo (QMC) [24, 25] at zero temperature. The experiments are generally compatible with most these calculations and confirm universal behavior [13, 26] in the unitarity limit as well as superfluidity at high temperatures [19, 22, 26, 27, 28].

The purpose of this work is to calculate sound velocities in the superfluid phase and demonstrate that first and second sound undergoes avoided crossing at the BEC-BCS crossover in uniform systems of strongly interacting Fermi atoms. Second sound reveals the presence of a superfluid and its critical temperature. The sound modes provide details about the equation of state at the BEC-BCS crossover and constrain crossover models.

The two-body interaction range is assumed to be short

as compared to the s-wave scattering length a and the interparticle spacing or k_F^{-1} , where $n = k_F^3/3\pi^2$ is the density of the Fermi atoms with two equally populated spin states. The physics can be expressed in terms the variable $x = 1/(ak_F)$, which varies from $-\infty$ in the dilute BCS limit through the unitarity limit $x = 0$ at the Feshbach resonance to $x \rightarrow +\infty$ in the molecular BEC limit.

The sound modes may be measured in experiments as density waves in the expanding clouds. Density or thermal perturbations in the trapped clouds may be introduced in the center of the cloud, for example, by a pulsed laser before expansion, or by other means [29]. It is important that the two sound modes mix as will shown below because both modes will be excited by either a density or thermal perturbation. As the sound modes travel through the cloud the density and temperature may vary. It may therefore be nontrivial to extract the sound speeds as function of density, temperature and scattering length, and useful to have a model calculation of both sound speeds.

The manuscript is organized such that the sound modes are first described in general, and then calculated in the BEC, BCS and unitarity limits respectively. The three limits are then approximately matched assuming a smooth crossover. Finally, summary and conclusions are given.

II. SOUND MODES

We shall in the following assume that the thermal excitations collide frequently and therefore can be assumed in local thermal equilibrium. This is the collisional limit where hydrodynamics apply. The transition from zero to first sound as systems change from collisionless to collisional is described in, e.g., [30] and the hydrodynamics of superfluids in [31, 32]. For a Fermi liquid the collision time $\tau \propto T^{-2}$ can become large at low temperatures. The hydrodynamic limit is therefore limited to long wavelength (or equivalently low frequency ω) sound modes such that $\tau\omega \ll 1$. This can always be achieved in a bulk system but not necessarily in a finite system where

*Also at Univ. of S.Denmark; Electronic address: hh@ddre.dk

collective frequencies ω are finite. However, it was found in [6, 7, 33] that axial and radial collective modes can be described by hydrodynamics at temperatures well below the superfluid transition. Thus atoms in traps appear collisional in a wide region around the unitarity limit. Yet the collisionless limit is expected in traps at very low temperatures [34].

The presence of a normal and superfluid components leads to two sound modes in the collisional limit referred to as first and second sound. Their velocities u_1 and u_2 are given by the positive and negative solutions respectively of [32, 35]

$$u^2 = \frac{c_S^2 + c_T^2}{2} \pm \sqrt{\left(\frac{c_S^2 + c_T^2}{2}\right)^2 - c_T^2 c_2^2}. \quad (1)$$

The thermodynamic quantities entering are the adiabatic $c_S^2 = (\partial P / \partial \rho)_S$, the isothermal $c_T^2 = (\partial P / \partial \rho)_T$, and the “thermal” $c_2^2 = \rho_s s^2 T / \rho_n c_V$ sound speed squared. The latter also acts as a coupling or mixing term. The difference between the adiabatic and isothermal sound speed squared can also be expressed as

$$c_S^2 - c_T^2 = \left(\frac{\partial s}{\partial \rho}\right)_T^2 \frac{\rho^2 T}{c_V}. \quad (2)$$

Here, $\rho = \rho_n + \rho_s = nm$ are the total, ρ_n the normal and ρ_s the superfluid mass densities, $s = S/\rho$ the entropy per unit mass and $c_V = T(\partial s / \partial T)_\rho$ the specific heat per unit mass.

The isothermal sound speed at zero temperature is in both the hydrodynamic limit and for a superfluid gas given by

$$\begin{aligned} c_{T=0}^2 &= \frac{n}{m} \left(\frac{\partial \mu}{\partial n} \right)_T \\ &= \frac{1}{3} v_F^2 \left[1 + \beta - \frac{3}{5} x \beta' + \frac{1}{10} x^2 \beta'' \right], \end{aligned} \quad (3)$$

where $v_F = \hbar k_F / m$, $\beta(x) = E_{int} / E_{kin}$ is the ratio of interaction to kinetic energy, $\beta' = d\beta/dx$, etc. QMC calculations [24, 25] find $\beta(x=0) = -0.57$ in the unitarity limit at zero temperature. The continuous crossover found in QMC and crossover models is confirmed experimentally for the pressure, chemical potential, β , collective modes [1, 3, 8] and recently also for the entropy [9].

In the dilute BCS limit and at low temperature $c_T = v_F \sqrt{(1 + (2/\pi)ak_F)/3}$. In the unitarity limit $c_T = v_F \sqrt{(1 + \beta(0))/3} \simeq 0.37v_F$. In the dilute BEC limit $c_T = \sqrt{(\pi/2)\hbar^2 na_M/m^2}$, where a_M is the molecular scattering length. According to QMC [25] and four-body calculations [36] $a_M \simeq 0.62a$, whereas in the Leggett and several other crossover models $a_M = 2a$.

We are mainly interested in the superfluid state at temperatures below the critical temperature, which generally is less than that in the BEC limit, $T \leq T_c \lesssim T_c^{BEC} = 0.218E_F$, the temperatures will be much smaller than the Fermi energy E_F (in units where $k_B = 1$). At such low




FIG. 1: First and second sound speeds for temperatures $T/T_c = 0.25, 0.5, 0.75$. The isothermal sound speed c_T is calculated from the Leggett crossover model. Sound speeds are calculated using entropies from Eq. (4) on the BEC side, Eq. (12) on the BCS side and from the measured entropy in the unitarity limit at $x = 0$ (see text).

temperature the isothermal sound speed is given by Eq. (3) in both the BCS and BEC limits. In the crossover model of Leggett the isothermal sound speed is also to a good approximation given by the zero temperature value for $T \lesssim T_c$. We shall therefore use Eq. (3) for c_T^2 in the following.

The entropy and mass densities have been calculated in the BEC [32, 35] and BCS [37] limits to leading order in ak_F . One should, however, be careful in extrapolating to the unitarity limit. For example a normal BEC is depleted as $a_M \rightarrow +\infty$ and eventually quenched [38] whereas in crossover models the condensate remains although it changes character from BEC to BCS superfluid. Likewise, the collective mode frequencies in trapped Fermi atoms increase for a normal BEC as the unitarity limit is approached [39]. Crossover models [33] and experiments [6, 8], however, find the opposite for a molecular BEC. The BEC and BCS limits will now be described followed up with a discussion on the extension and interpolation towards the unitarity limit in various crossover models.

III. BEC LIMIT

Crossover models and calculations find different molecular scattering length in the BEC limit as mentioned above. Also the effective mass M in the dispersion relation $E_q = q^2/2M$ differs between the molecular BEC limit where $M = 2m$ and the unitarity limit, where M becomes very large indicating that the dispersion relation is not quadratic but rather linear as a Bogoliubov phonon. For example, the pair susceptibility χ , which enters selfenergies, is calculated with different combinations of bare (G_0) and full (G) Green's functions. With $\chi = GG$ [17, 20] M increases monotonically as the uni-




FIG. 2: As Fig.1 but with c_T and $a_M = 0.62a$ from QMC. Furthermore, induced interactions are included on the BCS side which reduce T_c .

tarity limit is approached from the molecular BEC limit, whereas with $\chi = G_0 G_0$ [16, 18] M first decreases and then increases as $x \rightarrow 0_+$. With $\chi = G_0 G$ [19] M first increase, then decrease and finally increase again as $x \rightarrow 0_+$. The critical temperature has the opposite behavior as M because $T_c \propto 1/M$.

A normal BEC has a linear Bogoliubov dispersion relation $E_q = qc_T$ for temperatures below $T^* = 2\pi a_M n/M$ corresponding to $x \lesssim (a_M/a)(T_c/T)$. For $a_M = 0.62a$ and $T \sim T_c/2$ the quadratic dispersion relation is thus limited to $x \gtrsim 1$. Similar results are found in several crossover model [19, 23], where the molecular gas can on the BEC side ($x \gtrsim 1$) be described approximately by a dispersion relation $E_q = q^2/2M$, with an effective mass $M \simeq 2m$.

The corresponding entropy per particle is then given by the standard result

$$\frac{S}{N} = \sigma \frac{3}{2} \left(\frac{T}{T_c} \right)^{3/2}, \quad (4)$$

where $\sigma = 5\zeta(5/2)/3\zeta(3/2) = 0.856$ and the critical temperature is $T_c = 2\pi\hbar^2(n/2\zeta(3/2))^{2/3}/M = 0.218E_F(2m/M)$. The normal mass density is on the BEC side $\rho_n = \rho(T/T_c)^{3/2}$. From the entropy and mass densities on the BEC side we obtain for the mixing term $c_2^2 = \sigma(T/M)\rho_s/\rho$ and $c_S^2 - c_T^2 = \sigma(T/M)\rho_n/\rho$.

In the BEC limit we thus obtain the classical result for the first sound speed

$$u_1 = \sqrt{\sigma \frac{T}{M} + c_T^2}, \quad (5)$$

and the standard second sound speed $u_2^2 = c_T^2 \rho_s/\rho = 4\pi\hbar^2 a_M \rho_s/M^3$, or

$$u_2 = \sqrt{\frac{a_M k_F}{6\pi} [1 - (T/T_c)^{3/2}]} v_F. \quad (6)$$

As we approach the unitarity limit we observe that *avoided crossing* takes place between the two sound

modes caused by the mixing term in Eq. (1). The mixing acts similar to level repulsion between otherwise degenerate states in quantum mechanical systems. Instead of crossing the two sound speeds avoid crossing by an amount proportional to the mixing term, and first sound turns continuously into second sound and visa versa. As the avoided crossing occurs predominantly on the BEC side it can be described by using the BEC mass densities and entropies in Eq. (1). The resulting sound modes are shown in Fig. 1 with c_T and T_c from the Leggett crossover model which has $a_M = 2a$. For comparison the sound modes are calculated in Fig. 2 with c_T from QMC [25], which has $a_M = 0.62a$. In both cases the entropy of Eq. (4) with $M = 2m$ is assumed.

IV. BCS LIMIT

The thermodynamic quantities and sound speeds in BCS limit have been calculated in [37] from the mean field BCS equations of Leggett [15]

$$\frac{1}{g} = \sum_{\mathbf{k}} \left[\frac{1}{2E_k} - \frac{m}{\hbar^2 k^2} - \frac{f_k}{E_k} \right], \quad (7)$$

with coupling strength $g = -4\pi\hbar^2 a/m$. As usual $\varepsilon_k = \hbar^2 k^2/2m - \mu$, $E_k = \sqrt{\varepsilon_k^2 + \Delta^2}$ and $\Delta(T)$ the gap. The thermal distribution function is $f_k = (\exp(E_k/T) + 1)^{-1}$. With the equation for number density conservation

$$n = \sum_{\mathbf{k}} \left[1 - \frac{\varepsilon_k}{E_k} + 2 \frac{\varepsilon_k}{E_k} f_k \right], \quad (8)$$

the gap and chemical potential can be calculated as function of density, temperature and interaction strength. At zero temperature the last terms in Eqs. (7) and (8) vanish and the gap is

$$\Delta_0 = \left(\frac{8}{e^2} \right) E_F \exp(\pi/2ak_F). \quad (9)$$

Gorkov [40] included induced interactions, which reduce the gap to

$$\Delta_0 = (2/e)^{7/3} E_F \exp(\pi/2ak_F). \quad (10)$$

Induced interaction can be included in the gap equation (7) by replacing $a^{-1} \rightarrow a^{-1} - 2k_F \ln(4e)/3\pi$ on the left hand side. Hereby, not only the gap is corrected but also the thermodynamic potential [37]. In both cases $T_c = (\gamma/\pi)\Delta_0 = 0.567\Delta_0$.

The thermodynamic functions can be calculated from the thermodynamic potential per volume $\Omega = -P$. We make the standard assumption that the Hartree-Fock terms in the superfluid Ω_s and normal state Ω_n thermodynamic potentials are the same. The difference is then given in terms of the pairing coupling as

$$\Omega_s = \Omega_n + \int_0^\Delta d\Delta' \Delta'^2 \frac{d(1/g)}{d\Delta'}. \quad (11)$$

The first order temperature correction to the thermodynamic potential in the normal phase is $\Omega_n = -N(0)\pi^2 T^2/3$, where $N(0) = mk_F/2\pi^2$ is the level density.

Inserting the coupling of Eq. (7) into the thermodynamic potential (11) and again exploiting that $\Delta \ll \mu$, it reduces to

$$\Omega_s = -N(0) \left[\Delta^2 \left(\frac{1}{2} + \ln \frac{\Delta_0}{\Delta} \right) - 4T \int_0^\infty d\varepsilon_k \ln(1 - f_k) \right] \quad (12)$$

The crossover model thus arrives at the standard expression for Ω_s and therefore also the standard entropy density $S_s = -(\partial\Omega_s/\partial T)_{V,\mu}$, and specific heat $C_s = T(\partial S_s/\partial T)_{V,\mu}$ in superfluid phase. In the normal phase $S_n = C_n = N(0)2\pi^2 T/3$. Near T_c : $S_s/S_n(T) = 1 - (1 + \xi)(1 - T/T_c)$ and $C_s/C_n(T_c) = \xi - 3.77(1 - T/T_c)$. Here, $\xi = 1 + 12/7\zeta(3) \simeq 2.43$ is the ratio of superfluid specific heat just below T_c to the normal one just above T_c . At low temperatures $S_s = N(0)\sqrt{2\pi\Delta_0^3/T} \exp(-\Delta_0/T)$ and $C_s = N(0)\sqrt{2\pi\Delta_0^5/T^3} \exp(-\Delta_0/T)$.

Finally, we need the superfluid (London) density

$$n_s = n \left(1 + 2 \int_0^\infty d\varepsilon_k \frac{df_k}{dE_k} \right), \quad (13)$$

where $n = n_s + n_n$ is the total density. At low temperatures $n_s/n = 1 - \sqrt{2\pi\Delta_0/T} \exp(-\Delta_0/T)$ whereas $n_s/n = 2(1 - T/T_c)$ near T_c .

We now have all the thermodynamic quantities available for calculating the sound speeds in the BCS limit. Since thermal effects are small we find that $u_1 = c_S \simeq c_T$. The second sound is a pure thermal wave with velocity

$$u_2^2 \simeq c_2^2 = \frac{n_s}{n_n} \frac{S_s^2 T}{mn C_s}. \quad (14)$$

The dependence on temperature and x is shown in Fig. 3 as well as in Figs. 1 and 2 for three different temperatures.

At low temperatures the second sound is linear in temperature

$$u_2 = \frac{\sqrt{3}}{2} \frac{T}{E_F} v_F, \quad T \ll T_c. \quad (15)$$

Around $T \simeq 0.7T_c$ the second sound speed has a broad maximum of

$$u_2 \simeq 0.53 \frac{T}{E_F} v_F, \quad T \sim 0.7T_c. \quad (16)$$

Near the critical temperature $T_c - T \ll T_c$

$$u_2 = \frac{\pi}{\sqrt{\xi}} \frac{T}{E_F} \sqrt{1 - \frac{T}{T_c}} v_F. \quad (17)$$

The characteristic $u_2 \propto \sqrt{1 - T/T_c}$ behavior near T_c is the same as in the BEC limit, Eq. (6), but with a different prefactor.




FIG. 3: Thermodynamic quantities for a superfluid as calculated in the BCS limit vs. T/T_c . Shown are the gap $\Delta(T)/\Delta_0$, entropy $S_s(T)/S_n(T)$, specific heat $C_s(T)/C_n(T)\xi$, and superfluid density $n_s(T)/n$. The second sound is plotted in units of $v_F T_c/E_F$.

V. UNITARITY LIMIT AND CROSSOVER

At crossover the entropy receives contributions from thermal Fermi atoms and molecular bosons. The entropies and sound speeds reflect the underlying dispersion relation for low-lying excitations. On the BEC side for $T \gtrsim T^*$ the quadratic dispersion relation $E_q = \hbar^2 q^2/2M$ leads to an entropy $S \propto T^{3/2}$, whereas for $T \lesssim T^*$ the linear Bogoliubov phonon $E_q = \hbar q c_T$ yields $S \propto T^3$. On the BCS side the gap in the quasiparticle energy leads to a more complicated entropy as discussed above.

Generally, for a power law dispersion relation $E_q = q^\alpha$ the entropy and specific heat scale as $S \propto C_V \propto T^{3/\alpha}$ and the normal mass density $\rho_n = \rho(T/T_c)^{5/\alpha-1}$. Consequently $c_S^2 - c_T^2 \propto T^{3/\alpha+1}$ and $c_2^2 \propto (\rho_s/\rho) T^{2-2/\alpha}$. The resulting temperature dependence of the sound velocities follows from Eq. (1).

Model calculations [14, 15, 16, 17, 18, 19, 20, 21, 22, 23] find a smooth crossover of most thermodynamic quantities as function of density and scattering length at zero and finite temperature except at T_c , where superfluidity vanishes. We will therefore extrapolate the sound modes from the BCS and BEC limits towards crossover such that they are continuous around crossover and meet the sound speeds in the unitarity limit, which now will be calculated independently.

Experimentally, the entropy has recently been measured and found to scale as $S/N \simeq 20(T/T_F)^{1.53}$ in the unitarity limit at temperatures below $T_c \simeq 0.27 E_F$ [9]. In the unitarity limit the entropy is independent of scattering length due to universality [13] as other thermodynamic quantities [26]. Consequently, when $S/N \propto T^{3/2}$, the entropy density is independent of density and is therefore the same in bulk as in a trap at temperatures sufficiently low as compared to the Fermi temperature. Both

the measured T_c and entropy are then slightly larger in the unitarity limit than for the pure BEC of Eq. (4). With these adjustments of the parameters in Eq. (4), the sound velocities are calculated in the unitarity limit as in the BEC limit above as shown in Figs. 1 and 2 for $x = 0$. To guide the eye the unitarity limit point at $x = 0$ is connected smoothly in the crossover regime to the BEC limit, $x \geq 1$, and the BCS limit, $x \leq -1$.

The sound velocities in Figs. 1 and 2 are well behaved in the sense that both the BEC and BCS limits extrapolate continuously towards the unitarity limit. On the BCS side first sound is basically given by the isothermal sound speed, and the second sound increases exponentially with x as $T_c = (\gamma/\pi)\Delta$ as given by Eqs. (9) and (10). Around the unitarity limit their dependence on temperature and $x = 1/k_F a$ is more complicated because the two sound modes mix and the temperature dependence of the entropy is different from the BEC limit. On the BEC side the curves again extrapolates continuously towards the BEC limit, however, with avoided crossing of the two sound modes.

It should be noted that the curves in Figs. 1 and 2 are plotted for constant T/T_c . On the BEC side T_c is almost constant from the unitarity to the BEC limit, and the curves are therefore basically for constant temperature. On the BCS side, however, T_c decrease exponentially as $T_c \propto \exp(\pi x/2)$ (see Eqs.(9) and (10)), and therefore the temperatures also decrease. If the temperature were held fixed, superfluidity and the second sound would thus vanishes below a certain x where $T \geq T_c$.

The specific heat is a continuous function of temperature around T_c only in the BEC limit. It is discontinuous in the BCS limit by a factor $C_s/C_n = 2.43$. Recent calculations and experimental fits [19] at $x = 0.11$ find $C_s/C_n = 2.51 \pm 0.05$ at T_c . These experiments indicate that the discontinuity in the heat capacity has a maximum crossing over from the BCS value towards zero in the BEC limit. T_c also has a maximum at crossover in several crossover models [16, 18, 19].

The second sound speed squared generally vanishes with the superfluid density at the critical temperature,

i.e. $c_2 \propto \sqrt{T_c - T}$. This is observed in the BCS limit of Eq. (17)), in the BEC limit of Eq. (6) and also in the unitarity limit [13].

The sound speeds are continuous as function of x at crossover as long as T/T_c is fixed as shown in Figs. 1 and 2. As function of temperature first sound depends only little on temperature whereas second sound vanishes at T_c and the specific heat is discontinuous. Similar temperature dependency were found in Refs. [21, 41] at crossover.

VI. SUMMARY

The first and second sounds have been calculated in the molecular BEC, the BCS and the unitarity limits. It has been argued that the sound modes interpolate continuously between these limits, and that avoided crossing takes place on the BEC side due to mixing.

The mixing has the consequence that both sound modes can be excited and detected both as density and thermal waves in traps. In contrast the second sound is a pure thermal wave in the BCS and BEC Bogoliubov limits. In finite traps the sound waves could be excited in the center and propagate towards the surface so that the local density decreases. Consequently, $|x| = 1/|a|k_F$ increases and the sound velocities change as shown in Figs. 1 and 2.

The second sound mode reveals superfluidity and the critical temperature. Measurements of both sound modes constrain the equation of state, entropies, specific heat, etc. and thereby crossover models, which predict very different behavior of, e.g., the molecular scattering length in the BEC limit and effective molecular mass, critical temperature, etc. at crossover. The crossover and mixing of the sound modes will also be possible to study in mixtures of bosons and fermions. A number of different superfluids may also be created in optical lattices [42] (with corresponding Mott transitions) and possibly also supersolids.

-
- [1] K. M. O'Hara, S. L. Hemmer, M. E. Gehm, S. R. Granade, J. E. Thomas, *Science* **298** (2002) 2179. M. E. Gehm et al., *Phys. Rev. A* **68**, 011401 (2003); *cond-mat/0304633*.
 - [2] M. Greiner, C.A. Regal, D.S. Jin, *Nature* **426**, 537 (2003); *Phys. Rev. Lett.* **92**, 040403 (2004).
 - [3] T. Bourdel et al., *Phys. Rev. Lett.* **91**, 020402 (2003);
 - [4] M.W. Zwierlein et al., *Phys. Rev. Lett.* **91**, 250401 (2003); S. Gupta et al., *Science* **300**, 47 (2003).
 - [5] S. Jochim et al., *Phys. Rev. Lett.* **91**, 240402 (2003); M. Bartenstein et al., *ibid.* **92**, 120401 (2004)
 - [6] J. Kinast et al., *Phys. Rev. Lett.* **92**, 150402 (2004)
 - [7] M. Bartenstein et al., *Phys. Rev. Lett.* **92**, 203201 (2004)
 - [8] C. Chin, M. Bartenstein A. Altmeyer, S. Riedl, S. Jochim, J. Hecker Denschlag, R. Grimm, *Science* **305** 1128 (2004).
 - [9] J. Kinast et al., *cond-mat/0409283* See also, *Science* **307**, 1296 (2005).
 - [10] M. Greiner, C.A. Regal, J.T. Stewart, D.S. Jin, *cond-mat/0502411*.
 - [11] D. Vollhardt and P. Wölfle, *The Superfluid Phases of Helium 3*, Taylor and Francis, London 1990; *cond-mat/0012052*.
 - [12] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems*, McCraw-Hill 1971.
 - [13] T-L. Ho, *Phys. Rev. Lett.* **92**, 090402 (2004)
 - [14] D.M. Eagles, *Phys. Rev.* **186**, 456 (1969).
 - [15] A.J. Leggett, in *Modern Trends in the Theory of Condensed Matter*, ed. A. Pekalski and R. Przystawa, Lect. Notes in Physics Vol. 115 (Springer-Verlag, 1980), p. 13.

- [16] P. Nozières and S. Schmidt-Rink, J. Low Temp. Phys. **59**, 195 (1982).
- [17] R. Haussmann, Z. Phys. B**91**, 291 (1993).
- [18] C. A. R. Sá de Melo, M. Randeria, and J. R. Engelbrecht, Phys. Rev. Lett. **71**, 3202 (1993). M. Randeria, in “Bose-Einstein Condensation”, Ed. A. Griffin, D.W. Snoke, S. Stringari, Cambridge Univ. Press 1995, p. 355.
- [19] J. Stajic, J. N. Milstein, Qijin Chen, M. L. Chiofalo, M. J. Holland, K. Levin, Phys. Rev. A **69**, 063610 (2004).
- [20] A. Perali, P. Pieri, G.C. Strinati, Phys. Rev. Lett. **93**, 100404 (2004); *ibid.* **92**, 220404 (2004). P. Pieri, L. Pisani, G.C. Strinati, Phys. Rev. B **70**, 094508 (2004).
- [21] Y. Ohashi, A. Griffin, Phys. Rev. Lett. **89**, 130402 (2002); cond-mat/0410220; Phys. Rev. A **72**, 013601 (2005).
- [22] M. Holland et al., Phys. Rev. Lett. **87**, 120406 (2001).
- [23] L.M. Jensen, cond-mat/0412431.
- [24] J. Carlson, S-Y. Chang, V. R. Pandharipande, K. E. Schmidt, Phys. Rev. Letts. **91**, 50401 (2003). S-Y. Chang et al., Phys. Rev. A **70**, 043602 (2004)
- [25] G.E. Astrakharchik, J. Boronat, J. Casulleras, S. Giorgini, Phys. Rev. Lett. **93**, 200404 (2004)
- [26] H. Heiselberg, Phys. Rev. A **63**, 043606 (2001); J. Phys. B: At.Mol.Opt.Phys. **37**, 1 (2004).
- [27] E. Timmermans et al., Phys. Lett. A **285**, 228 (2001)
- [28] G.M. Falco and H.T.C. Stoof, cond-mat/0409621.
- [29] J. Thomas, priv. comm.
- [30] G. Baym and C.J. Pethick, *Landau Fermi-liquid Theory*, Wiley & Sons, 1991.
- [31] L.D. Landau and E.M. Lifshitz, *Fluid Mechanics*, Second edition, (Pergamon, New York, 1987).
- [32] C.J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, Cambridge Univ. Press, 2002.
- [33] H. Hu, A. Minguzzi, X-J Liu, M.P. Tosi, Phys. Rev. Lett. **93**, 190403 (2004). H. Heiselberg, Phys. Rev. Lett. **93**, 040402 (2004).
- [34] G. M. Bruun and H. Smith, cond-mat/0504734
- [35] A. Griffin and E. Zaremba, Phys. Rev. A **56**, 4839 (1997).
- [36] D.S. Petrov, C. Salomon, and G.V. Shlyapnikov, cond-mat/0309010.
- [37] H. Heiselberg, cond-mat/0409077.
- [38] S. Cowell et al., Phys. Rev. Lett. **88**, 210403 (2002).
- [39] S. Stringari, Europhys. Lett. **65**, 749 (2004).
- [40] L. P. Gorkov & T. K. Melik-Barkhudarov, JETP **13**, 1018 (1961); H. Heiselberg, C. J. Pethick, H. Smith and L. Viverit, Phys. Rev. Letts. **85**, 2418 (2000).
- [41] T. Domanski & J. Ranninger, Phys. Rev. B **70**, 184513 (2004).
- [42] D.-W. Wang, M.D. Lukin, E. Demmler, cond-mat/0410494.